

# Genus-One String Amplitudes from Conformal Field Theory

Luis Fernando Alday

University of Oxford

Current Developments in QFT and Gravity - Kolkata


# What will this talk be about?

Conformal Field Theory techniques to attack Scattering Amplitudes in theories of Gravity/String Theory.

Based on work with A. Bissi, E. Perlmutter.

# Scattering amplitudes

Scattering Amplitudes: One of the most important observables in any quantum field theory.



- $A(g, s, t, u)$  depends on many things:
  - Which particles you are scattering (their masses, charges, etc)
  - The parameters of your theory  $g$ .
  - The momenta of the particles being scattered:

$$s = -(p_1 + p_2)^2, \quad t = -(p_1 - p_3)^2, \quad u = -(p_1 - p_4)^2$$

- They can teach us a lot about a theory!

# Scattering amplitudes in pure gravity


## General Relativity: Einstein Hilbert Lagrangian

$$\mathcal{L}_{EH}[g] = \frac{1}{G_N} \sqrt{-g} \mathcal{R}$$

Consider a scattering process:  $g_{\mu\nu} = \eta_{\mu\nu} + \sqrt{G_N} h_{\mu\nu}$

$$\mathcal{L}_{EH} = (\partial h)^2 + \sqrt{G_N} h (\partial h)^2 + \dots$$

Scattering of gravitons:




- UV divergences: the gravitons get too close to each other!

# String theory and UV divergences


- Replace gravitons by closed strings of finite length  $\sqrt{\alpha'}$

Point particles



VS

Strings



- At low energies,  $p^2 \ll 1/\alpha'$  we recover GR (or SUGRA).


GR as an effective field theory

$$\mathcal{L} = (\partial h)^2 + \sqrt{G_N} \left( h(\partial h)^2 + \alpha'^{1/2} h^2 (\partial h)^2 + \alpha' h (\partial h)^3 + \dots \right)$$


- At high energies stringy corrections give a UV completion of GR!

# String theory scattering amplitudes

- In string theory we have two parameters




- We would like to compute scattering amplitudes



$$= A(g_s, \alpha', s, t, u)$$

# String theory scattering amplitudes

- The computation organises in a genus expansion



$$A^{(\text{genus } 0)}(\alpha', s, t, u) + g_s^2 A^{(\text{genus } 1)}(\alpha', s, t, u) + g_s^4 A^{(\text{genus } 2)}(\alpha', s, t, u) + \dots$$


At genus zero and in flat space: Virasoro-Shapiro amplitude

$$A^{(\text{genus } 0)}(\alpha', s, t, u) = \frac{\Gamma(-\frac{\alpha' s}{4})\Gamma(-\frac{\alpha' t}{4})\Gamma(-\frac{\alpha' u}{4})}{\Gamma(1 + \frac{\alpha' s}{4})\Gamma(1 + \frac{\alpha' t}{4})\Gamma(1 + \frac{\alpha' u}{4})}$$

- Even in flat space higher genus terms are notoriously hard to compute! For curved space time, we have not idea how to do this!
- We will use an alternative approach.

## AdS/CFT duality

String theory on  $AdS_5 \times S^5$   $\Leftrightarrow$  Quantum field theory living in the 4d boundary of  $AdS_5$



- $\mathcal{N} = 4$  Super conformal Yang-Mills, with  $SU(N)$  gauge group.
- Most symmetric 4d theory!

Parameters:  $g_s$  and  $R$ .


Parameters:  $g_{YM}$  and  $N$ .

## Dictionary

$$g_s = \left( \text{Diagram of a genus-1 surface} \right) \approx \frac{1}{N}, \quad \frac{R^2}{\alpha'} = \sqrt{g_{YM}^2 N} \equiv \sqrt{\lambda}$$



String amplitudes on  $AdS_5 \times S^5 \leftrightarrow$  correlators of local operators!



$$A(g_s, \frac{\alpha'}{R^2}, s, t, u) \leftrightarrow \langle \mathcal{O}(x_1) \mathcal{O}(x_2) \mathcal{O}(x_3) \mathcal{O}(x_4) \rangle$$

## String Amplitudes on $AdS_5 \times S^5$

## Correlators in $\mathcal{N} = 4$ SYM

Genus expansion

$1/N$  expansion

Stringy corrections to SUGRA

$1/\lambda$  corrections

Graviton on  $AdS$

$\mathcal{O}_2$ : protected scalar of dim. 2  
in the stress-tensor multiplet

KK modes on  $S^5$

$\mathcal{O}_p$ : protected ops of dim.  $p$

- Compute  $\langle \mathcal{O}_2(x_1)\mathcal{O}_2(x_2)\mathcal{O}_2(x_3)\mathcal{O}_2(x_4) \rangle$  in a  $1/N$  and  $1/\lambda$  expansion.
- Recently made possible thanks to the analytic bootstrap!

# Analytic bootstrap

Four-point function of identical operators:

$$\langle \mathcal{O}(x_1)\mathcal{O}(x_2)\mathcal{O}(x_3)\mathcal{O}(x_4) \rangle = \frac{\mathcal{G}(u, v)}{x_{12}^{2\Delta_{\mathcal{O}}} x_{34}^{2\Delta_{\mathcal{O}}}}$$

$$\text{where } u = \frac{x_{12}^2 x_{34}^2}{x_{13}^2 x_{24}^2}, v = \frac{x_{14}^2 x_{23}^2}{x_{13}^2 x_{24}^2}$$

## Conformal partial wave decomposition

- OPE:  $\mathcal{O} \times \mathcal{O} = \sum_i \mathcal{O}_i + \text{descendants}$


$$\mathcal{G}(u, v) = \sum_{\mathcal{O}} \begin{array}{c} 2 \\ \diagdown \\ \text{---} \\ \diagup \\ 1 \end{array} \begin{array}{c} \mathcal{O} \\ \text{---} \\ c_{\phi\phi\mathcal{O}} \end{array} \begin{array}{c} \mathcal{O} \\ \text{---} \\ c_{\phi\phi\mathcal{O}} \end{array} \begin{array}{c} 3 \\ \diagup \\ \text{---} \\ \diagdown \\ 4 \end{array}$$

$$\mathcal{G}(u, v) = \sum_{\mathcal{O}_{\Delta, \ell}} \underbrace{c_{\Delta, \ell}^2}_{\text{OPEcoef.}} \underbrace{g_{\Delta, \ell}(u, v)}_{\text{conformal blocks}}$$

# Analytic bootstrap

## Crossing symmetry

$$v^{\Delta \circ} \mathcal{G}(u, v) \underbrace{=}_{x_1 \leftrightarrow x_3} u^{\Delta \circ} \mathcal{G}(v, u)$$




Highly non-trivial equation!

$$v^{\Delta \circ} \sum_{\Delta, l} c_{\Delta, l}^2 g_{\Delta, l}(u, v) = u^{\Delta \circ} \sum_{\Delta, l} c_{\Delta, l}^2 g_{\Delta, l}(v, u)$$

# Numerical vs Analytic bootstrap

Study this equation in different regions,  $u = z\bar{z}$ ,  $v = (1 - z)(1 - \bar{z})$




- In the Euclidean regime  $\bar{z} = z^*$ .
- We can study crossing around  $u = v = \frac{1}{4}$
- Starting point of the numerical bootstrap.

- In the Lorentzian regime  $z, \bar{z}$  are independent real variables and we can consider  $u, v \rightarrow 0$ .
- Starting point of the analytic (light-cone) bootstrap!

# Analytic bootstrap

- In Minkowski space we can have  $x_{23}^2 \rightarrow 0, x_{23} \neq 0$  (small  $v$ , any  $u$ )
- When some operators become null-separated the correlator develops singularities:



- The whole correlator can be reconstructed from these singularities!

## LSPT/Inversion formula

$$\mathcal{G}(u, v) = \int dudv K(u, v) \text{sing}[\mathcal{G}(u, v)] + \text{ambiguities}$$

- Hence the spectrum and OPE coef. of intermediate operators.

# Example: Generalised free fields

- Simplest solution: - Generalised free fields (CFTs at  $N = \infty$ )

$$\mathcal{G}^{(0)}(u, v) = 1 + \left(\frac{u}{v}\right)^{\Delta_{\mathcal{O}}} + u^{\Delta_{\mathcal{O}}}$$

## Intermediate operators

$$1 + \underbrace{\left(\frac{u}{v}\right)^{\Delta_{\mathcal{O}}} + u^{\Delta_{\mathcal{O}}}}_{\text{Double trace operators}}$$

Identity operator

$$\mathbb{I} \quad \mathcal{O} \square^n \partial_{\mu_1} \cdots \partial_{\mu_\ell} \mathcal{O}$$
$$\Delta = \ell = 0 \quad \Delta_{n,\ell} = 2\Delta_{\mathcal{O}} + 2n + \ell$$

- Let's include  $1/N^2$  corrections to this!

# Large $N$ CFTs

## AdS/CFT

Large  $N$  CFT in  $D$ -dimensions  
(GFF + corrections)

$\Leftrightarrow$

Gravitational theory in  
 $AdS_{D+1}$

$\frac{1}{N^2}$  expansion in CFT  $\leftrightarrow$  genus/loops in  $AdS$

$$\mathcal{G} = \underbrace{\text{Diagram 1}}_{N^0} + \underbrace{\text{Diagram 2} + \text{Diagram 3}}_{1/N^2} + \underbrace{\text{Diagram 4} + \text{Diagram 5}}_{1/N^4} + \dots$$

- Direct computations in  $AdS$  are hard...Let's use CFT!



$$\mathcal{G}(u, v) = \mathcal{G}^{(0)}(u, v) + \boxed{\frac{1}{N^2} \mathcal{G}^{(1)}(u, v)} + \frac{1}{N^4} \mathcal{G}^{(2)}(u, v) + \dots$$

## Two Sources of corrections

- 1 Double trace operators will acquire corrections:

$$\begin{aligned}\Delta_{n,\ell} - \ell &= 2\Delta_{\mathcal{O}} + 2n + \frac{1}{N^2} \gamma_{n,\ell}^{(1)} + \dots \\ a_{n,\ell} &= a_{n,\ell}^{(0)} + \frac{1}{N^2} a_{n,\ell}^{(1)} + \dots\end{aligned}$$

- 2 We can also have new intermediate operators at order  $1/N^2$ .

Which corrections are consistent with crossing symmetry and the structure of (null) singularities?

# Large $N$ holographic CFTs

## Exchange solutions

In large- $N$  CFT's dual to gravitational theories we always have the stress tensor! (dual to the graviton)

$$\mathcal{O} \times \mathcal{O} = 1 + [\mathcal{O}, \mathcal{O}]_{n,\ell} + \frac{1}{N^2} T_{\mu\nu}$$

- This produces a well defined divergence in the null-limit:

$$\mathcal{G}^{(1)}(u, v) \sim \frac{h_T(u)}{v^{\Delta_{\mathcal{O}} - (d-2)/2}}$$

- From the divergences we can fix  $\gamma_{n,\ell}^{(1)}, a_{n,\ell}^{(1)}$ !
- This solution corresponds to  $AdS$  graviton exchange

$$\mathcal{G}_{grav}^{(1)}(u, v) \sim \text{Diagram}$$

## Truncated solutions

- In addition we can have special solutions consistent with no extra null divergence (corresponding to the ambiguities).
- $\gamma_{n,\ell}^{(1)}, a_{n,\ell}^{(1)}$  vanish to all orders in  $1/\ell!$
- But! we can have solutions with finite support in the spin.
- Truncated solutions  $\leftrightarrow$  local interactions in the  $AdS$  bulk.

$$\mathcal{G}_{trunc}^{(1)}(u, v) \sim \text{Diagram}$$

# Large $N$ holographic CFTs

$$\mathcal{G}(u, v) = \mathcal{G}^{(0)}(u, v) + \frac{1}{N^2} \mathcal{G}^{(1)}(u, v) + \boxed{\frac{1}{N^4} \mathcal{G}^{(2)}(u, v)} + \dots$$

- Null divergences at order  $1/N^4$  arise from the square of the anomalous dimension:


$$\mathcal{G}^{(2)}(u, v) \sim \log^2 v \sum_{n, \ell} \left( \gamma_{n, \ell}^{(1)} \right)^2 \partial_n^2 g_{n, \ell}(v, u)$$

- Given  $\left( \gamma_{n, \ell}^{(1)} \right)^2$  we can reconstruct  $\mathcal{G}^{(2)}(u, v)$ !

$$\left( \gamma^{(1)} \right)^2 = \left( \gamma_{grav}^{(1)} + \gamma_{trunc}^{(1)} \right)^2 = \gamma_{grav}^{(1)} \times \gamma_{grav}^{(1)} + \gamma_{grav}^{(1)} \times \gamma_{trunc}^{(1)} + \gamma_{trunc}^{(1)} \times \gamma_{trunc}^{(1)}$$

leads to the following contributions


# Contributions to order $1/N^4$



# Double expansion in $\mathcal{N} = 4$ SYM

$\mathcal{N} = 4$  SYM falls into this category! with the following structure

$$\mathcal{G}^{(0)}(u, v) + \frac{1}{N^2} \left( \mathcal{G}_{grav}^{(1)}(u, v) + \frac{1}{\lambda^{3/2}} \mathcal{G}_{st}^{(1)}(u, v) + \dots \right) + \frac{1}{N^4} \left( \mathcal{G}_{grav}^{(2)}(u, v) + \frac{1}{\lambda^{3/2}} \mathcal{G}_{st}^{(2)}(u, v) + \dots \right)$$



We can compute all these contributions!

The string amplitude in  $AdS_5 \times S^5$  is  $M(s, t, u)$

$$\mathcal{G}(U, V) = \int_{-i\infty}^{i\infty} ds dt U^s V^t M(s, t, u) \Gamma^2(2-s) \Gamma^2(2-t) \Gamma^2(2-u)$$

where  $s + t + u = 2$

- Crossing symmetry  $\rightarrow M(s, t, u)$  is completely symmetric.
- Gravity solution  $\rightarrow M(s, t, u)$  a meromorphic function:

$$M_{sugra}^{(1)}(s, t, u) = \frac{1}{(s-1)(t-1)(u-1)}$$


- Truncated solutions/stringy corrections  $\rightarrow M(s, t, u)$  is a polynomial. Nice basis

$$\sigma_2 = s^2 + t^2 + u^2, \quad \sigma_3 = s^3 + t^3 + u^3$$

# Mellin space

Genus zero expansion:

$$M^{\text{genus } 0}(s, t, u) = \frac{1}{(s-1)(t-1)(u-1)} + \alpha'^3 a + \alpha'^5 (b\sigma_2 + b_1) + \alpha'^6 (c\sigma_3 + c_1\sigma_2 + c_2) + \dots$$



*gravity*       $\mathcal{R}^4$        $\partial^4 \mathcal{R}^4$        $\partial^6 \mathcal{R}^4$

Flat space limit:  $s, t, u \rightarrow \infty, \alpha' \rightarrow 0$  with  $s\alpha', t\alpha', u\alpha' \sim \text{fixed}$

## Genus 0 flat space limit

$$M^{\text{genus } 0}(s, t, u) \rightarrow \frac{1}{stu} + \alpha'^3 a + \alpha'^5 b\sigma_2 + \alpha'^6 c\sigma_3 = \text{VS amplitude}$$

- $b_1, c_1, c_2$  are curvature corrections.



# Mellin space and flat space limit

- $M^{\text{genus } 1}(s, t, u)$  is a mess but we are able to compute it!

Genus-one string amplitude on  $AdS_5 \times S^5$  in a  $\alpha'$  expansion

$$M_{\text{grav.}}^{\text{genus } 1}(s, t, u) + \alpha'^3 M_{\text{st},1}^{\text{genus } 1}(s, t, u) + \alpha'^5 M_{\text{st},2}^{\text{genus } 1}(s, t, u) + \dots$$

- In the flat space limit they simplify a lot!

$$M_{\text{grav.}}^{\text{genus } 1}(s, t, u) \rightarrow \text{Box function in 10D}$$

$$M_{\text{st},1}^{\text{genus } 1}(s, t, u) \rightarrow s^4 \log s + t^4 \log t + u^4 \log u$$

$$\begin{aligned} M_{\text{st},2}^{\text{genus } 1}(s, t, u) \rightarrow & (87s^6 + s^4(t-u)^2) \log s \\ & + (87t^6 + t^4(u-s)^2) \log t \\ & + (87u^6 + u^4(s-t)^2) \log u \end{aligned}$$

- Agrees exactly with the known results! **[Green, Russo, Vanhove]**

# Conclusions

- We have developed an efficient machinery to study  $1/N$  corrections to holographic correlators (in various dimensions).
- We have provided the first genus one computation of a string theory amplitude in curved space-time.
- We can answer detailed questions, such as the structure of UV divergences, and in the flat space limit we recover the known structure.
- Can we now start asking quantitative questions about quantum gravity/string theory on curved spaces?