

Senior Professor of Theoretical Chemistry, School of Chemical Sciences, Indian Association for the Cultivation of Science, and Adjunct Research Professor, Indian Institute of Science Education and Research Kolkata (IISER K) 07TH February, 2023, 4.00 pm at FERMION Hall

ABSTRACT :: The first principle based Beyond Born-Oppenheimer (BBO) theory [1-3] is presented for the construction of highly accurate potentially coupled matrix (diabatic) to study multi-mode multi-surface photoelectron spectra of NO 3 - and 1,3,5-C 6 H 6 F 3 + [4-5], reactive scattering processes of H 3 + and F + H 2 [6-9], and phase transition phenomena of orthorhombic manganites [10]. Jahn-Teller (JT) and Renner-Teller (RT) types of conical intersections (CIs) along with Pseudo Jahn-Teller (PJT) interactions in those systems and semicircular CI seam (rather than a CI point) between the ground and excited state are the new interesting observations. [4-6, 9] The theoretically calculated spectra of the titled systems show good peak by peak correspondence with the experimental and other theoretical findings. [4-5] Reaction attributes (reaction probabilities, cross-sections and/or rate constants) of D + +H 2 and H+H 2 + processes obtained from scattering dynamics over global diabatic surfaces of H 3 + system [7,8] in hyperspherical coordinates for total angular momentum zero and non-zero situations exhibit good accord with the experimentally measured ones. On the other hand, reactive scattering calculation over the diabatic surfaces of F+H 2 system (incorporating spin-orbit couplings) is quite encouraging. Optical spectra of REMnO 3 (RE = La, Pr, Nd, Sm, Eu, Gd, Tb, Dy) shows anomalous temperature dependence around the Neel temperature. This behaviour could be due to Jahn-Teller effect both in ground and excited state. Such observations have been investigated [10,11] theoretically to interpret the experimental spectra due to the excitations of the quantum rotors.

References:

- [1] B. Mukherjee, K. Naskar, S. Mukherjee, S. Ghosh, T. Sahoo and S. Adhikari, Int. Rev. Phys. Chem. 38, 287 (2019).
- [2] K. Naskar, S. Mukherjee, B. Mukherjee, S. Ravi, S. Mukherjee, S. Sardar and S. Adhikari, J. Chem. Theory Comput. 16, 1666-1680 (2020).
- [3] J. Dutta, S. Mukherjee, K. Naskar, S. Ghosh, B. Mukherjee, S. Ravi and S. Adhikari, Phys. Chem. Chem. Phys. (Perspective) 22, 27496-27524 (2020).
- [4] B. Mukherjee, S. Mukherjee, S. Sardar, K. R. Shamasundar and S. Adhikari, Chem. Phys. 515, 350 (2018).
- [5] S. Mukherjee, S Ravi, J. Dutta, S. Sardar and S. Adhikari, Phys. Chem. Chem. Phys. 24, 2185 (2022).
- [6] S. Mukherjee, D. Mukhopadhyay and S. Adhikari, J. Chem. Phys., 141, 204306 (2014).
- [7] S. Ghosh, S. Mukherjee, B. Mukherjee, S. Mandal, R. Sharma, P. Chaudhury, and S. Adhikari, J. Chem. Phys. 147, 074105 (2017).
- [8] S Ghosh, T Sahoo, M Baer and S Adhikari, J. Phys. Chem. A 125, 731-745 (2021).
- [9] B. Mukherjee, K. Naskar, S. Mukherjee, S. Řavi, K. R. Shamasundar, D. Mukhopadhyay and S. Adhikari, J. Chem. Phys. 153, 174301 (2020).
- [10] J. Dutta, S. Ravi, S. Mukherjee, A. K. Ojha and S. Adhikari, J. Phys. Chem. A, 126, 691 (2022).
- [11] J. Dutta, N. N. Kovaleva and S. Adhikari, J. Chem. Phys., 150, 064703 (2019).

Webinar link

YouTube link